

PS03lite Orchidopexy for a Palpable Testicle (Child)

Expires end of December 2019

A more detailed version of this procedure specific information sheet is available. Please ask your health practitioner at Joondalup Health Campus for a copy.

Write questions or notes here:



**HEALTH CONSUMERS'
COUNCIL**
YOUR VOICE ON HEALTH

This document will give you information about an orchidopexy. If you have any questions, you should ask your GP or other relevant health professional.

What is an orchidopexy?

An orchidopexy is an operation to bring a testicle down into the scrotum. The testicles develop in a baby boy's abdomen when he is in the womb. The testicles usually move down into the scrotum by 35 weeks of pregnancy. Sometimes a testicle does not come down normally (see figure 1).




Figure 1

Both testicles usually move down into the scrotum

What are the benefits of surgery?

Surgery should prevent your child from having serious complications. Your child's fertility should improve, and he will find it easier when he is an adult to examine his testicles to check for any problems.

Are there any alternatives to surgery?

If a testicle has not reached the scrotum by the age of six months, it is unlikely to do so without surgery.

What does the operation involve?

The operation is performed under a general anaesthetic and usually takes 45 minutes to an hour.

Your surgeon will usually perform the operation through a cut on the groin and a small cut on the scrotum. They will free up the testicle and bring it down into the scrotum.

If your surgeon finds a small testicle that is unlikely to function, they will usually remove it.

What complications can happen?

1 General complications of any operation

- Pain
- Infection of the surgical site (wound)
- Bleeding
- Unsightly scarring

2 Specific complications of this operation

- Developing a collection of blood or fluid
- Shrinking of the testicle
- Prevention of sperm passing to the penis
- The testicle may return to its original position
- Reduction in fertility of a testicle that is brought down

How soon will my child recover?

Your child should be able to go home the same day.

It is usual for children to return to school after about a week.

Your child should not play sports or ride a bicycle for 6 weeks.

Summary

An orchidopexy is an operation to bring a testicle down into the scrotum. If left untreated, serious complications can happen.

Acknowledgements

Author: Mr Shailinder Singh DM FRCS (Paed. Surg.), Mr Gregor Knepil FRCS (Ed.) and Mr Jonathan Sutcliffe FRCS

Illustrations: Medical Illustration Copyright © Medical-Artist.com

This document is intended for information purposes only and should not replace advice that your relevant health professional would give you.